

Clinico-epidemiological Profile of Sexually Transmitted Infections in a Tertiary Hospital: A Cross-Sectional Study

Alina Karki¹, Varsha Manandhar², Rupak Maharjan¹

¹ Department of Dermatology, Nepal Armed Police Force Hospital, Balambu, Kathmandu, Nepal

² Department of Community Medicine, Manipal College of Medical Sciences, Fulbari, Pokhara, Nepal

Corresponding Author:

Dr. Alina Karki

Department of Dermatology
Nepal Armed Police Force Hospital
Balambu, Kathmandu, Nepal.
Email: alinakarki1701@gmail.com

Article History:

Received date: July 17, 2025

Revised date: August 01, 2025

Accepted date: August 07, 2025

Published date: August 28, 2025

Online Access



DOI:10.64772/mjapfn118

Abstract

Introduction: Sexually transmitted infections are communicable diseases that occur due to sexual contact. They are a major global health problem that affects the quality of life of patients. Knowledge of the frequency and pattern of sexually transmitted infections helps in early diagnosis and appropriate treatment. The aim of the study was to determine the pattern of various sexually transmitted infections in a tertiary hospital in Kathmandu.

Methods: This cross-sectional study was conducted among patients visiting the Dermatology outpatient department of a tertiary hospital in Kathmandu from 1st Dec 2019 to 30th Nov 2023 after ethical approval from the Nepal Health Research Council (Reference Number: 681/2023). The data were retrieved from the records of the department, entered and analyzed using Statistical Package for the Social Sciences version 21.

Results: A total of 165 patients were included in the study, among which 121(73.33%) were male and 44(26.67%) were female, with a male-to-female ratio of 2.7:1. Most of the patients were married and in the age group 26 to 35 years. Most common sexually transmitted infections were condyloma acuminata 56(33.95%), urethral discharge syndrome 37(22.43%), herpes genitalis 28(16.97%), vaginal discharge syndrome 20(12.12%), syphilis 18(10.90%), molluscum contagiosum 3(1.81%), and others 3(1.81%).

Conclusions: Viral infections were the most common sexually transmitted infections in our study.

Keywords: *condyloma acuminata; epidemiology; sexually transmitted infections.*

Introduction

Sexually Transmitted Infections (STIs) are a group of communicable diseases spread predominantly through sexual contact and caused by over 30 pathogens.¹ Most are asymptomatic, stigmatized, and slowly growing into hidden epidemics.² Globally, STIs remain a major public health problem.³ According to the World Health Organization (WHO) data of 2020, globally more than one million people

acquired curable STIs every day of which chlamydia, gonorrhea, syphilis and trichomoniasis were the most common.⁴ Untreated STIs like gonorrhea and syphilis can cause infertility, adverse pregnancy outcomes and Human Immunodeficiency Virus (HIV) co-infection.⁵⁻⁷ Hospital-based studies of STI. Nepal lacks national STI surveillance, though hospital-based studies report a prevalence of 0.3%-1.2%.^{8,9}

How to cite (Vancouver Style)

Karki A, Manandhar V, Maharjan R. Clinico-epidemiological profile of Sexually Transmitted Infections in a Tertiary Hospital: A Cross- Sectional Study. Med J APF Nepal. 2025;1(1):32-36.

© The Author(s) 2025 This work is licensed under a Creative Commons Attribution 4.0 International License. (CC BY-NC)



This study aims to describe the clinico-epidemiological profile of STIs in patients visiting the outpatient department of the dermatology department of Nepal Armed Police Force (APF) Hospital.

Methods

This descriptive cross-sectional study was conducted after ethical approval from the Nepal Health Research Council (Reference Number: 681/2023). All eligible cases, above 15 years of age, irrespective of gender, diagnosed with STIs who visited the dermatology OPD of Nepal APF Hospital, Kathmandu from 1st December 2019 to 30th November 2023, were included in the study. HIV infection cases were excluded from the study. The data were retrieved from the database from the department. Only the details of patients that were mentioned under the diagnosis of sexually transmitted infections were collected based on the clinical and laboratory evaluation. This information was entered in proforma and verified by principal investigator. The data were entered and analyzed in Statistical Package for the Social Sciences (SPSS) version 21. Mean and standard deviation for continuous data, frequencies, and proportions for categorical variables were calculated.

Results

A total of 12,191 patients visited the dermatology OPD during the four-year study period, out of which 165 (1.35%) patients were diagnosed with STIs. Among them, there were 121 (73.33%) males and 44 (26.67%) females, with male to female ratio of 2.7:1. Majority of patients (76, 46.06%) belonged to age group 26-35 years. Patients ranged from 17 years to 68 years of age, with mean age of 32.70 ± 10.09 years (Figure 1).




Figure 1: Age and sex-wise distribution of the patients with STIs

In our study 119 (72.12%) patients were married, 39 (23.64%) were unmarried, five (3.03%) were divorced and two (1.21%) were widowed. Among the cases with STI, 102 (61.81%) had multiple sex partners.

Most cases 163 (98.87%) had heterosexual mode of contact, while two cases (1.13%) had history of homosexual contact.

The most common STIs diagnosed were condyloma acuminata, seen in 56 (33.95%) cases, followed by Urethral Discharge Syndrome seen in 37 (22.43%) cases. The frequencies and pattern of STIs observed in our study are shown in Table 1.

Table 1: Frequency and Pattern of Sexually Transmitted Infections (n=165).

Diagnosis	Male(%)	Female(%)	Total(%)
Condyloma acuminata	47(38.85)	9(20.45)	56(33.95)
Urethral Discharge Syndrome	37(30.58)	-	37(22.43)
Herpes Genitalis	20(16.52)	8(18.18)	28(16.97)
Vaginal Discharge Syndrome	-	20(45.45)	20(12.12)
Syphilis	13(10.75)	5(11.36)	18(10.90)
Molluscum contagiosum	2(1.65)	1(2.28)	3(1.81)
Others	2(1.65)	1(2.28)	3(1.81)
Total	121(100)	44(100)	165(100)

Discussion

Out of 12,191 patients, who visited Dermatology OPD of Nepal APF Hospital, 165 (1.35%) patients were diagnosed with STIs. Data on prevalence of STI is scarce in Nepal, however hospital based studies done has shown STIs in 0.3%-1.2% of Dermatology OPD visits.^{8,9} Similarly, a hospital based study from Northeast India had shown STIs on 0.80% cases.¹⁰ However, a study on pregnant women at antenatal clinic in Nepal had shown prevalence of any STIs as 8.6% (7.1% of trichomoniasis, 1.5 % of chlamydia trachomatis).¹¹ This difference in prevalence of STIs could be attributed to variation in study site, populations, social and cultural differences, stigma and low OPD visits at COVID era.

In our study STIs were seen more in males, in consensus with other hospital-based studies in Nepal.^{5,12-14} Few studies from India have showed male predominance while another by Thapar et al. showed more STI cases in female.¹⁵⁻¹⁷ This was due to more ANC referral in STI clinic in their study.¹⁷ Male predominance in our study might be due to military personnel being primary beneficiary in our hospital. Less females with STIs could be secondary to social restrictions, stigma and reluctance. Also, inability of our study to include patients from gynecology department could have resulted in less female cases.

Sexually active age group 15-45 years constituted 149 (90.30%) of study population, similar to study done by Paudel et al. with 90.8% cases in 15-49 years age group.⁶ Majority 76 (46.06%) of our patients were of age group 26-35 years. Study done by Banger et al. had majority (54.32%) in 21-30 years age group and Vora et al. had 59.70% cases in 25-44 years age group.^{18,19} This is also similar to other studies done in Nepal.^{13,14} This age group being most sexually active, may have more sexual partners and concurrent partnerships resulting in increased incidence of the disease.

STIs were more common in married patients, similar to other studies from Nepal and abroad.^{5,13,16,19} Married people tend to use less protection measures and may take privilege of marital status to have hidden sexual contacts. Partners of case may be the cause for transmission. However some studies show more STIs in unmarried subjects.^{12,14} This finding may be due to majority of cases in early twenties in these studies. Heterosexual contact 163 (98.87%) was most common mode of sexual contact similar to multiple studies.¹⁸⁻²¹ Homosexual contact was seen 1.13% cases similar to studies by Banger et al. and Devi et al. who showed in 1.28% and 1.2% cases respectively.^{18,21}

Most common STIs in our study was condyloma acuminata seen in 56 (33.95%) cases, similar to findings by Paudel et al. and Gyawalee et al.¹²⁻¹³ Herpes infections were seen in 28 (16.97%) cases which is similar to study done by Narayanan et al. where 17.9% of cases had this infection.²⁰ It was the third most common STIs in our study. It was the most common STIs in other studies.^{18,19} It is also the foremost cause of STI according to World Health Organization.²² The high incidence of condyloma and herpes is in accordance with other studies.^{12,13,16} Worldwide trend of decreasing bacterial sexual infections and increasing trend of viral infections which may also be due to tendency of viral infections to be latent and persistent.¹⁸

Urethral Discharge Syndrome (UDS) was present in 37 (22.43%) and Vaginal Discharge Syndrome (VDS) in 20 (12.12%) cases. UDS was the second most common STIs in our study. In study done by Karki et al. it was seen in 20.1% and 17% of cases respectively.⁵ A study done by Sharma et al. had VDS as most common STIs.²³ It was due to increased referral to STI clinic from gynecology OPD. Low VDS findings in our study may be due to more females visiting gynecology department and seeking treating prior to visiting dermatology department in our hospital.

Syphilis cases were less 18 (10.90%) in our study. However, it was the most common findings in various studies done in Nepal and India.^{5,9,10,20} These studies attributed it to routine ANC screening in

hospitals and screening of migrant workers for job opportunities. The less incidence in our study might be due to less screening for travels, job opportunities and decreased OPD visits due to travel restrictions as most of our study period is during COVID times. It might also be due to self-medications, antibiotic usage during this time which may have interfered with diagnosis.

Conclusions

In our study, condyloma acuminata was the most common STIs. High prevalence of STIs in married cases highlights importance of contact tracing and partner treatment.

Source of Funding: None

Acknowledgement

We would like to acknowledge staffs of Dermatology OPD of Nepal APF hospital for helping in data collection and maintaining records.

Conflict of Interest: None

Author's Contribution and ORCID iDs

Dr. Alina Karki

Conceptualization, Design, Literature review, Data analysis, Drafting.

ID : <https://orcid.org/0009-0000-8991-7189>

Dr. Varsha Manandhar

Literature review, Data analysis, Manuscript editing and review

ID : <https://orcid.org/0000-0003-3788-3794>

Dr. Rupak Maharjan

Manuscript editing and review

ID : <https://orcid.org/0009-0008-6307-6519>

References

1. Deng M, Chen J, Wang Z, Zheng R, Pang W, Sun R, et al. Trends in the Incidence of Common Sexually Transmitted Infections at the Global, Regional and National Levels, 1990–2021: Results of the Global Burden of Disease 2021 Study. *Trop Med Health*. 2025 May 16;53(1):70. DOI: [10.1186/s41182-025-00744-2](https://doi.org/10.1186/s41182-025-00744-2)
2. John Kurien A, Gupta B. A Clinico-epidemiological Study of Non-Retroviral Sexually Transmitted Diseases in Patients Attending Dermatology Department in a Tertiary Care Hospital. *Indian J Clin Exp Dermatol*. 2021 Feb 28;7(1):30–4. DOI: [10.18231/j.ijced.2021.006](https://doi.org/10.18231/j.ijced.2021.006)

3. Huang J, Lin S, Pan J, Lu L, Liu B, Zheng J, et al. Global Burden of Sexually Transmitted Infections: 1990 - 2019. In Review. 2021 Aug. DOI: [10.21203/rs.3.rs-847312/v1](https://doi.org/10.21203/rs.3.rs-847312/v1)
4. World Health Organisation. Sexually Transmitted Infections (STIs). 2025 Jul 28. Available from: [https://www.who.int/news-room/fact-sheets/detail/sexually-transmitted-infections-\(stis\)](https://www.who.int/news-room/fact-sheets/detail/sexually-transmitted-infections-(stis))
5. Karki A, Shrestha R, Parajuli N. Pattern of Sexually Transmitted Infections in a Tertiary Care Hospital: A Five-Year Retrospective Study. *Grande Med J*. 2019 Dec 31;1(2):84–7. DOI: [10.3126/gmj.v1i2.27058](https://doi.org/10.3126/gmj.v1i2.27058)
6. World Health Organisation. Guidelines for the Management of Symptomatic Sexually Transmitted Infections. 2025 Jul 28. Available from: <https://iris.who.int/bitstream/handle/10665/342523/9789240024168-eng.pdf>
7. Wu MY, Gong HZ, Hu KR, Zheng HY, Wan X, Li J. Effect of Syphilis Infection on HIV Acquisition: A Systematic Review and Meta-Analysis. *Sex Transm Infect*. 2021 Nov;97(7):525–33. DOI: [10.1136/sextrans-2020-054706](https://doi.org/10.1136/sextrans-2020-054706)
8. Thapa DP, Rana A. Awareness and Pattern of Sexually Transmitted Diseases: A Hospital Based Study. *Nepal J Dermatol Venereol Leprol*. 2023 Oct 2;21(2):20–7. DOI: [10.3126/njdv.v21i2.55298](https://doi.org/10.3126/njdv.v21i2.55298)
9. Paudel U, Sharma A, Chapagain L, Gartaula M, Gupta P, Parajuli S. Patterns of Sexually Transmitted Infections in a Tertiary Care Hospital: An Alarming Rise in the Prevalence of Syphilis. *Nepal Med Coll J*. 2025 Apr 17;27(1):65–9. DOI: [10.3126/nmcj.v27i1.77545](https://doi.org/10.3126/nmcj.v27i1.77545)
10. Gupta B, Datta S, Das K, Roy J. The Trend of STD in COVID Times- A Retrospective Hospital-Based Study from Northeast India. *IP Indian J Clin Exp Dermatol*. 2022;8(1):43–6. DOI: [10.18231/j.ijced.2022.009](https://doi.org/10.18231/j.ijced.2022.009)
11. Dev R, Adhikari SP, Dongol A, Madhup SK, Pradhan P, Shakya S, et al. Prevalence Assessment of Sexually Transmitted Infections Among Pregnant Women Visiting an Antenatal Care Center of Nepal: Pilot of the World Health Organization's Standard Protocol for Conducting STI Prevalence Surveys Among Pregnant Women. *PLoS One*. 2021;16(4):e0250361. DOI: [10.1371/journal.pone.0250361](https://doi.org/10.1371/journal.pone.0250361)
12. Paudel S, Parajuli N, Dahal S, Paudel S. Sexually Transmitted Infections in A Tertiary Hospital of Kathmandu: A Retrospective Study. *Nepal J Dermatol Venereol Leprol*. 2021 Oct 4;19(2):44–8. DOI: [10.3126/njdv.v19i2.37686](https://doi.org/10.3126/njdv.v19i2.37686)
13. Gyawalee M, Pokhrel DB. Pattern of Sexually Transmitted Infections and Sexual Behavior in Patients with Genital Symptoms. *Nepal J Dermatol Venereol Leprol*. 2014;12(1):20–7. DOI: [10.3126/njdv.v12i1.10592](https://doi.org/10.3126/njdv.v12i1.10592)
14. Karn D, Amatya A, Aryal ER, Ks S, Timalisina M. Prevalence of Sexually Transmitted Infections in a Tertiary Care Centre. *Kathmandu Univ Med J*. 2011;9(34):44–8. DOI: [10.3126/kumj.v9i2.6287](https://doi.org/10.3126/kumj.v9i2.6287)
15. Sarkar S, Shrimal A, Das J, Choudhury S. Pattern of Sexually Transmitted Infections: A Profile from a Sexually Transmitted Infections Clinic of a Tertiary Care Hospital of Eastern India. *Ann Med Health Sci Res*. 2013 Apr;3(2):206–9. DOI: [10.4103/2141-9248.113663](https://doi.org/10.4103/2141-9248.113663)
16. Rajiv S, Hashba B. A Five-Year Retrospective Study of Pattern of Sexually Transmitted Diseases in a Tertiary Care Hospital in North Kerala. *Int J Contemp Med Res*. 2017;4(3):614–7. DOI: [10.21276/ijcmr.2018.5.5.21](https://doi.org/10.21276/ijcmr.2018.5.5.21)
17. Thapar R, Riyaz M, Kaur N. Prevalence and Pattern of Sexually Transmitted Diseases in a Tertiary Care Hospital in Chamba, Himachal Pradesh. *Int J Contemp Med Res*. 2018;5(5):E5–9. DOI: [10.21276/ijcmr.2018.5.5.21](https://doi.org/10.21276/ijcmr.2018.5.5.21)
18. Banger HS, Sethi A, Malhotra S, Malhotra SK, Kaur T. Clinico-Epidemiological Profile of Patients Attending Suraksha Clinic of Tertiary Care Hospital of North India. *Indian J Sex Transm Dis AIDS*. 2017;38(1):54–9. DOI: [10.4103/2589-0557.203436](https://doi.org/10.4103/2589-0557.203436)
19. Vora R, Anjaneyan G, Doctor C, Gupta R. Clinico-Epidemiological Study of Sexually Transmitted Infections in Males at a Rural-Based Tertiary Care Center. *Indian J Sex Transm Dis AIDS*. 2011 Jul 1;32(2):86–9. DOI: [10.4103/0253-7184.85410](https://doi.org/10.4103/0253-7184.85410)
20. Narayanan B. A Retrospective Study of the Pattern of Sexually Transmitted Diseases During a Ten-Year Period. *Indian J Dermatol Venereol Leprol*. 2005;71(5):333–7. DOI: [10.4103/0378-6323.19996782](https://doi.org/10.4103/0378-6323.19996782)
21. Devi SA, Vetrichevvel TP, Pise GA, Thappa DM. Pattern of Sexually Transmitted Infections in a Tertiary Care Centre at Puducherry. *Indian J Dermatol*. 2009;54(4):347–9. DOI: [10.4103/0019-5154.57611](https://doi.org/10.4103/0019-5154.57611)
22. Implementing the Global Health Sector Strategies on HIV, Viral Hepatitis and Sexually Transmitted Infections, 2022–2030. 2024. Available from: <https://www.who.int/publications/i/item/9789240094925>

23. Sharma A, Rattan R, Sood A. Pattern of Sexually Transmitted Infections in a District Hospital from Himachal Pradesh. *Int J Community Med Public Health*. 2017 Mar 28;4(4):1028–31. DOI: [10.18203/2394-6040.ijcmph20171319](https://doi.org/10.18203/2394-6040.ijcmph20171319)

Bios

Dr. Alina Karki is an Assistant Professor and Consultant Dermatologist at Department of Dermatology, Nepal Armed Police Force Hospital, Balambu, Kathmandu, Nepal. Her expertise combines academic teaching with clinical practice, focusing on skin health and advanced dermatological treatments in Nepal. Her areas of interest are Tropical Dermatology, Allergology, Laser and Aesthetics.

Email: alinakarki1701@gmail.com

Dr. Varsha Manandhar is an Assistant Professor at Manipal College of Medical Sciences, Pokhara. She holds MD degree in Community Medicine and Tropical Diseases. Her Areas of research are Non-communicable disease, Infant and young children feeding and self-medication.

Email: varshamanandhar42@gmail.com

Dr. Rupak Maharjan is Consultant Dermatologist. He serves as Head of Department of Dermatology at Nepal Armed Police Force Hospital, Balambu, and Kathmandu, Nepal. He has over a decade of experience in clinical dermatology. His area of research includes Clinical Dermatology, Tropical diseases and Infectious diseases.

Email: rupakmaharjan@yahoo.com